

## Follow-up

The Doctor will usually want to check that there are no problems a few weeks after fitting an IUD. After this, there is no need for any routine check until it is time to remove the IUD. However, see your Doctor at any time if you have any problems or queries. Most women have no problems and the IUD can remain in place for several years.

It can be removed at any time by a trained Doctor or Nurse. Fertility returns as soon as it is removed. If you plan to have it removed, but do not want to get pregnant, then use other methods of contraception (such as condoms) from seven days before it is removed. This is because sperm can last up to seven days after having sex.

You can use sanitary towels or tampons for periods when an IUS is in place.

You should consult your Doctor if any of the following occur:

- Prolonged abdominal pain after an IUS is inserted.
- A delayed period, or bleeding between periods.
- A delayed period and lower abdominal pain (which may be due to an ectopic pregnancy)
- Vaginal discharge with or without pain (which may indicate infection).
- If you suspect that the IUD has come out or is coming out. It is usually possible to feel the threads of the IUD inside the vagina to check it is in place. If you cannot feel the threads then use other contraceptive methods (such as condoms) until you have been checked by a Doctor or Nurse.



**CUMBERLAND HOUSE**  
**PRIMARY HEALTH CARE CENTRE**

Cumberland House

Waters Green Medical Centre  
Sunderland Street  
Macclesfield  
Cheshire  
SK11 6JL

Phone: 01625 428081  
Fax: 01625 503128  
[www.cumberlandhousesurgery.nhs.uk](http://www.cumberlandhousesurgery.nhs.uk)

**Intrauterine  
Device  
(IUD)  
Copper Coil**

## What is the Intrauterine Device?

An Intrauterine Device (IUD) is a small device made from plastic and copper, and is otherwise known as "the coil". It can be replaced quite easily into the uterus (womb) by a trained Doctor.

## How does the IUD work as a contraceptive?

It works mainly by stopping the egg and sperm from meeting. It may also prevent the fertilised egg from attaching to the lining of the uterus. The copper also has a spermicidal effect (kills sperm).

## How effective is the IUD for contraception?

Modern IUDs are more than 99% effective. This means that up to 1 woman in 100 will become pregnant each year using this method of contraception. (Compare this to when no contraception is used. More than 80 in 100 sexually active women who do not use contraception become pregnant within one year). There are many different types of IUD. They can stay in place for between 5-10 years, depending on the type of IUD inserted.

## What are the advantages of the IUD?

Once an IUD is inserted you can forget about any contraception. So, unlike the pill, you don't need to think about contraception every day. It does not interfere with sex. It is not a hormonal method so it has no side-effects on the rest of the body. This means that it will not affect your mood, weight or libido. Having an IUD does not increase your risk of having any type of cancer in the future. Most women can have an IUD if they wish.

## What are the disadvantages of the IUD?

Although the majority of women with an IUD have no problems, the following may occasionally occur:

### HEAVY, PAINFUL PERIODS

Some women find that their periods become heavier, longer or more painful with an IUD. This means that the IUD may not be suitable if you already have heavy or painful periods. There is a special IUD called the intrauterine system (IUS), which is like an IUD, but it also releases a progestogen hormone into the uterus.

This is an effective treatment for heavy periods as well as a contraceptive. (See leaflet "Intrauterine System" - IUS). However, painful and heavy periods can still be treated in the same way as in women who do not have an IUD. For example, by taking anti-inflammatory painkillers or other medicines during your periods.

### INFECTION

There is a small risk of an infection of the uterus (pelvic infection) when you have the IUD fitted. Infection is not caused by the IUD. It is caused by sexually transmitted infections already present, usually within the first 20 days after insertion. Therefore, a check for infection of the vagina or cervix may be offered by taking a swab before an IUD is inserted. You should not have an IUD inserted if you have an infection which has not been treated.

### ECTOPIC PREGNANCY

The chance of becoming pregnant is very small if you use an IUS. However, if you do become pregnant, there is a slightly increased risk of having an ectopic pregnancy. This means the pregnancy is in the fallopian tube and not in the uterus. This is rare, but serious. See a Doctor urgently if you miss a period and develop some lower abdominal pain.

### EXPULSION

Rarely the IUD may come out without you noticing.

### DAMAGE TO THE WOMB

The fitting of an IUD can very rarely cause damage to the uterus (womb).

## How is the IUD fitted?

This is usually done towards the end of a period or shortly after.

However, it can be fitted at any time provided that you are certain you are not pregnant.

You will need to have a vaginal examination. The Doctor will pass a small instrument into your uterus to check its size and position. An IUD is then fitted. You will be taught how to feel the threads of the IUD so you can check it is in place. It is best to check the threads regularly eg once a month just after a period.

Fitting an IUD can sometimes be uncomfortable. Once the IUD has been inserted some women have crampy pains like period pains for a few hours afterwards. These can be eased by painkillers such as paracetamol. Light vaginal bleeding may also occur for a short while.

